arXiv:0811.2394v1 [hep-ph] 14 Nov 2008

LPT-08-90
CPHT-PC086.1108

$\gamma^* \to \rho_T$ impact factor with twist three accuracy

I.V. Anikin*, D.Yu. Ivanov†, B. Pire**, L. Szymanowski‡ and S. Wallon§

*Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia
†Sobolev Institute of Mathematics and Novosibirsk State University, 630090 Novosibirsk, Russia
**Centre de Physique Théorique, École Polytechnique, CNRS, 91128 Palaiseau, France
‡Soltan Institute for Nuclear Studies, Warsaw, Poland
§LPT, Université d'Orsay, CNRS, 91404 Orsay, France

Abstract. We evaluate the impact factor of the transition $\gamma^* \to \rho_T$ taking into account the twist 3 contributions. We show that a gauge invariant expression is obtained with the help of QCD equations of motion. Our results are free of end-point singularities. This opens the way to a consistent treatment of factorization for exclusive processes with a transversally polarized vector meson.

Keywords: impact factor, diffraction, QCD
PACS: 12.38.Bx, 13.60.Le

MOTIVATION

The study of exclusive reactions in the generalized Bjorken regime has been the scene of significant progresses in the recent years, thanks to the factorization properties of the leading twist amplitudes [1] for deeply virtual Compton scattering and deep exclusive meson production. It however turned out that transversally polarized rho meson production did not enter the leading twist controlable case [2] but only the twist 3 more intricate part of the amplitude [3, 4]. An understanding of the quark-gluon structure of a transversally polarized vector meson is however an important task of hadronic physics if one cares about studying confinement dynamics. This quark gluon structure may be described by distribution amplitudes which have been discussed in great detail [5]. On the experimental side, a continuous effort has been devoted to the exploration of $\pi\pi$ photo and electro-production, from moderate to very large energy. The kinematical analysis of the final meson pair allows then to separate the different helicity amplitudes, hence to measure the transversally polarized ρ meson production amplitude. Although non-dominant for deep electroproduction, this amplitude is by no means negligible and needs to be understood in terms of QCD. Up to now, experimental information comes from electroproduction on a proton or nucleus. Future progress may come from real or virtual photon photon collisions, which may be accessible either at electron positron colliders or in ultraperipheral collisions at hadronic colliders, as recently discussed [6, 7].

CALCULATION OF THE IMPACT FACTOR

We are interested in the description of high energy reactions such as

$$\gamma^*(q) + \gamma^*(q') \to \rho_T(p_1) + \rho(p_2) \tag{1}$$

or

$$\gamma^*(q) + N \to \rho_T(p_1) + N \tag{2}$$

where the virtual photons carry large squared momenta $q^2 = -Q^2$ ($q'^2 = -Q'^2$) $\gg \Lambda^2_{QCD}$, and the Mandelstam variable s obeys the condition $s \gg Q^2,\, Q'^2, -t \simeq \underline{r}^2$. Neglecting meson masses, one considers for reaction (1) the light cone vectors p_1 and p_2 as the vector meson momenta ($2\,p_1 \cdot p_2 = s$); the "plus" light cone direction is directed along p_1 and the “minus” light cone direction along p_2 with vector n defined as $p_2/(p_1 \cdot p_2)$. In this Sudakov light-cone basis, transverse euclidian momenta are denoted with underlined letters. The virtual photon momentum q reads $q = p_1 - \frac{Q^2}{2}\,n$. The impact representation of the scattering amplitude for the reaction (1) is

$$\mathscr{M} = \frac{is}{(2\pi)^2} \int \frac{d^2\underline{k}}{\underline{k}^2} \Phi_1^{ab}(\underline{k}, \underline{r} - \underline{k}) \int \frac{d^2\underline{k}'}{\underline{k}'^2} \Phi_2^{ab}(-\underline{k}', -\underline{r} + \underline{k}') \int\limits_{\delta - i\infty}^{\delta + i\infty} \frac{d\omega}{2\pi i} \left(\frac{s}{s_0}\right)^{\omega} G_\omega(\underline{k}, \underline{k}', \underline{r}) \tag{3}$$

where G_ω is the 4-gluons Green function which obeys the BFKL equation. G_ω reduces to $1/\omega\,\delta(\underline{k} - \underline{k}')\underline{k}^2/(\underline{r} - \underline{k})^2$ within Born approximation. We focus here on the $\gamma^* \to \rho$ impact factor Φ of the subprocess

$$g(k_1, \varepsilon_1) + \gamma^*(q) \to g(k_2, \varepsilon_2) + \rho_T(p_1)\,, \tag{4}$$

which is the integral of the κ-channel discontinuity of the S-matrix element $\mathscr{S}_\mu^{\gamma_T^*\, g \to \rho_T\, g}$:

$$\Phi^{\gamma^* \to \rho}(\underline{k}, \underline{r} - \underline{k}) = e^{\gamma^* \mu}\, \frac{1}{2s} \int \frac{d\kappa}{2\pi} \, \mathrm{Disc}_\kappa \, \mathscr{S}_\mu^{\gamma^*\, g \to \rho\, g}(\underline{k}, \underline{r} - \underline{k})\,, \tag{5}$$

where $\kappa = (q + k_1)^2$ denotes the Mandelstam variable s for the subprocess (4). Note that the two reggeized gluons have so-called non-sense polarizations $\varepsilon_1 = \varepsilon_2^* = p_2\sqrt{2/s}$. Considering now the forward limit for simplicity, the gluon momenta read

$$k_1 = \frac{\kappa + Q^2 + \underline{k}^2}{s} p_2 + k_T\,, \quad k_2 = \frac{\kappa + \underline{k}^2}{s} p_2 + k_T\,, \quad k_1^2 = k_2^2 = k_T^2 = -\underline{k}^2\,. \tag{6}$$

The impact factor Φ can be calculated within the collinear factorization. It is a convolution of perturbatively calculable hard-scattering amplitudes H and soft correlators S involving relevant ρ-meson distribution amplitudes, symbolically written as

$$\Phi = \int d^4 l \cdots \mathrm{tr}[H(l \cdots) S(l \cdots)]\,, \tag{7}$$

integrating over the n ($n \geq 2$) body phase-space. Working within the light-cone collinear factorization framework, let us first derive the contribution of the diagrams with the quark-antiquark correlators. The basic tool is to decompose any hard coefficient function $H(l)$ around a dominant “plus" direction:

$$H(l) = H(xp) + \left.\frac{\partial H(l)}{\partial l_\alpha}\right|_{l = xp} (l - xp)_\alpha + \ldots\,, \qquad (l - xp)_\alpha \approx l^T_\alpha, \tag{8}$$

where the first term is associated with the usual twist-2 contribution while the term with the derivative with the kinematical twist 3. After Fourier transform, this second hard term is accompanied in (7) by a soft correlator involving derivatives of the fields. These contributions are shown on Fig.1a (no transverse derivatives), and Fig.1b (with transverse derivatives).



FIGURE 1. Contributions to the impact factor with the quark-antiquark correlators without transverse derivatives (a), with transverse derivatives (b), and with the quark-antiquark-gluon correlators (c, d).

This allows us to factorize the amplitude in the momentum space. After a Fierz decomposition of the Dirac matrices $\psi(0)\,\bar\psi(z)$ and $\psi(0)\,i\overleftrightarrow{\partial_\perp}\,\bar\psi(z)$, the amplitude takes, up to higher twist terms, the simple factorized form symbolically written as:

$$\Phi \sim \int dx\,\left\{\mathrm{tr}\left[H_{q\bar q}(xp)\,\Gamma\right]\,S^{\Gamma}_{q\bar q}(x) + \mathrm{tr}\left[\partial_\perp H_{q\bar q}(xp)\,\Gamma\right]\,\partial_\perp S^{\Gamma}_{q\bar q}(x)\right\} \tag{9}$$

where Γ stands for some γ-matrix from a full set and the soft part reads symbolically

$$\begin{aligned} S^{\Gamma}_{q\bar q}(x) &= \int \frac{d\lambda}{2\pi}\,e^{-i\lambda x}\langle\rho(p)|\bar\psi(\lambda n)\,\Gamma\,\psi(0)|0\rangle \\ \partial_\perp S^{\Gamma}_{q\bar q}(x) &= \int \frac{d\lambda}{2\pi}\,e^{-i\lambda x}\langle\rho(p)|\bar\psi(\lambda n)\,\Gamma\, i\overleftrightarrow{\partial_\perp}\,\psi(0)|0\rangle\,. \end{aligned} \tag{10}$$

Within twist 3 order, the correlators which need to be parameterized are thus[1] [4]

$$\langle\rho(p)|\bar\psi(z)\gamma_\mu\psi(0)|0\rangle \overset{\mathscr{F}_1}{=} m_\rho\, f_\rho\,[\varphi_1(x)\,(e^*\cdot n)p_\mu + \varphi_3(x)\,e^{*T}_\mu]\,, \tag{11}$$

$$\langle\rho(p)|\bar\psi(z)\gamma_5\gamma_\mu\psi(0)|0\rangle \overset{\mathscr{F}_1}{=} m_\rho\, f_\rho\, i\varphi_A(x)\,\varepsilon_{\mu\lambda\beta\delta}\,e^{*T}_\lambda\, p_\beta\, n_\delta \tag{12}$$

and, for the quark-antiquark operators with the transverse derivatives,

$$\langle\rho(p)|\bar\psi(z)\gamma_\mu i\overleftrightarrow{\partial^T_\alpha}\,\psi(0)|0\rangle \overset{\mathscr{F}_1}{=} m_\rho\, f_\rho\,\varphi^T_1(x)\,p_\mu e^{*T}_\alpha\,, \tag{13}$$

$$\langle\rho(p)|\bar\psi(z)\gamma_5\gamma_\mu i\overleftrightarrow{\partial^T_\alpha}\,\psi(0)|0\rangle \overset{\mathscr{F}_1}{=} m_\rho\, f_\rho\, i\,\varphi^T_A(x)\,p_\mu\,\varepsilon_{\alpha\lambda\beta\delta}\,e^{*T}_\lambda\,p_\beta\, n_\delta\,. \tag{14}$$

A careful treatment of the correlators needs to take into account the equations of motion for the fermion field which read $\langle i\overrightarrow{\hat D}(0)\psi(0)\,\bar\psi(z)\rangle = 0$, and the corresponding one for the antifermion field, where $i\overrightarrow{D}_\mu = i\overrightarrow{\partial}_\mu + gA_\mu$. This shows that one must also consider diagrams such as the one shown on Fig. 1c, d, with the three body quark-antiquark-gluon nonlocal operators. One parameterizes these new non-perturbative objects as

---

[1] In the following definitions, $\mathscr{F}_1$ and $\mathscr{F}_2$ denote 2 and 3-body relative Fourier transforms along p.

$$\begin{aligned}
&\langle \rho(p)|\bar{\psi}(z_1)\gamma_\mu g A^T_\alpha(z_2)\psi(0)|0\rangle \overset{\mathcal{F}_2}{=} m_\rho\, f_3\, B(x_1,x_2)\, p_\mu e^{*T}_\alpha\,,\\
&\langle \rho(p)|\bar{\psi}(z_1)\gamma_5\gamma_\mu g A^T_\alpha(z_2)\psi(0)|0\rangle \overset{\mathcal{F}_2}{=} m_\rho\, f_3\, i D(x_1,x_2)\, p_\mu \varepsilon_{\alpha\lambda\beta\delta}\, e^{*T}_\lambda\, p_\beta\, n_\delta\,, \qquad (15)
\end{aligned}$$

where fractions x_1, $\bar{x_2}$ and $x_2 - x_1$ correspond to the quark, antiquark and gluon. We decompose the impact factor as the sum of non-flip (Φ_{nf}) and flip (Φ_f) contributions

$$\Phi^{\gamma^*_T\to\rho_T}(\underline{k}^2) = \Phi^{\gamma^*_T\to\rho_T}_{nf}(\underline{k}^2)\,(-e_{\gamma^*}\cdot e^*) + \Phi^{\gamma^*_T\to\rho_T}_{f}(\underline{k}^2)\left[\frac{(e_{\gamma^*}\cdot k)(e^*\cdot k)}{\underline{k}^2} + \frac{(e_{\gamma^*}\cdot e^*)}{2}\right]. \qquad (16)$$

The gauge invariance of the total amplitude is guaranteed if $\Phi(\underline{k}^2 = 0) = 0$. This is satisfied when using the equations of motion. The procedure is quite intricate and will be discussed in detail in a forthcoming publication. We stress that the impact factor is gauge-invariant *provided* the three body correlator contribution is taken into account. We only display the results obtained in the Wandzura-Wilczek approximation

$$\Phi^{\gamma^*_T\to\rho_T}_{nf\,(WW)}(\underline{k}^2) = -\frac{e\, m_\rho\, f_\rho}{\sqrt{2}\,Q^2}\,\frac{\delta^{ab}}{2N_c}\int\limits_0^1 dx\,\frac{\underline{k}^2\left(\underline{k}^2 + 2\,Q^2 x\bar{x}\right)}{\left(\underline{k}^2 + Q^2 x\bar{x}\right)^2}\left[\int\limits_0^x dv\frac{\varphi_1(v)}{\bar{v}} + \int\limits_x^1 dv\frac{\varphi_1(v)}{v}\right], \qquad (17)$$

$$\Phi^{\gamma^*_T\to\rho_T}_{f\,(WW)}(\underline{k}^2) = \frac{e\, m_\rho\, f_\rho}{\sqrt{2}\,Q^2}\,\frac{\delta^{ab}}{2N_c}\int\limits_0^1 dx\,\frac{2\,\underline{k}^2\, Q^2}{\left(\underline{k}^2 + Q^2 x\bar{x}\right)^2}\left[\bar{x}^2\int\limits_0^x dv\frac{\varphi_1(v)}{\bar{v}} + x^2\int\limits_x^1 dv\frac{\varphi_1(v)}{v}\right], \qquad (18)$$

where $\bar{x} = 1 - x$. Our results are free of end point singularities, thanks to the presence of the transverse momentum $\underline{k}$ in the denominators which regulates them.

ACKNOWLEDGMENTS

This work is partly supported by the ECO-NET program, contract 18853PJ, the French-Polish scientific agreement Polonium, the grant ANR-06-JCJC-0084, the RFBR (grants 06-02-16215, 08-02-00334, 08-02-00896) and the Polish Grant N202 249235.

REFERENCES

1. J. C. Collins, L. Frankfurt, M. Strikman, Phys. Rev. D **56**, 2982 (1997).
2. M. Diehl *et. al.* Phys. Rev. D **59**, 034023 (1999); J. C. Collins *et.al.*, Phys. Rev. D **61**, 114015 (2000).
3. L. Mankiewicz and G. Piller, Phys. Rev. D **61**, 074013 (2000).
4. I. V. Anikin and O. V. Teryaev, Phys. Lett. B **554**, 51 (2003), Nucl. Phys. A **711**, 199 (2002), Phys. Lett. B **509**, 95 (2001); I. V. Anikin *et. al.* Phys. Rev. D **62**, 071501 (2000).
5. P. Ball *et. al.* Nucl. Phys. B **529**, 323 (1998) and Phys. Rev. D **54**, 2182 (1996).
6. D. Yu. Ivanov *et. al.* Eur. Phys. J. C **38**, 195 (2004), Nucl. Phys. B **732**, 183 (2006) and Eur. Phys. J. C **49**, 947 (2007).
7. B. Pire *et. al.* Eur. Phys. J. C **44**, 545 (2005); Phys. Lett. B **639**, 642 (2006); R. Enberg *et. al.* Eur. Phys. J. C **45**, 759 (2006), Erratum-ibid. C **51**, 1015 (2007); M. Segond *et. al.* Eur. Phys. J. C **52**, 93 (2007); B. Pire *et. al.* arXiv:0810.3817.